



Keeping Your Balance

Module 6

Cognitive Strategies for Preventing Mania

Cognitive Changes in Mania	2
The Cognitive Behavioural Approach to Preventing Mania	4
An Example of a “Balance” Sheet	5
The “Balance” Sheet	6
Module Summary	7

The information provided in this document is for information purposes only. Please refer to the full disclaimer and copyright statements available at www.cci.health.gov.au regarding the information on this website before making use of such information.

COGNITIVE CHANGES IN MANIA



The onset of mania or hypomania can often be a pleasant experience for people with bipolar disorder. Positive changes in mood can be quite dramatic, and people often feel energised, excited, and optimistic. On the other hand people sometimes find that mania and hypomania begin with irritability and agitation, or that they experience both feelings of dysphoria and euphoria at the beginning of a manic episode. For those who experience mixed episodes, quick changes can occur between being in a good mood and extreme irritability.

The symptoms of mania, hypomania, or mixed episodes usually emerge over a period of days to weeks. Often people become aware of a pattern in the development of their symptoms, for example, hypomania or mania may begin with one symptom such as insomnia, and then progress to other symptoms (e.g., increased sexual interest, feelings of euphoria). Changes in the nature of people's thoughts, or preoccupation with certain ideas can also be part of this progression. Some people learn to recognise these changes in their thinking and can say, "It's happening again. I always start thinking that way when I'm getting high." This recognition can be a critical step for intervening early to prevent a full-blown manic episode.



Cognitive Symptoms of Mania & Hypomania

The cognitive changes associated with mania or hypomania include changes in the way people or situations are viewed, and changes in the amount and quality of new ideas. These changes can act as a signal that a manic episode is developing, and that action needs to be taken to prevent the symptoms from getting worse.

Increased Optimism & Grandiosity

When people are manic or hypomanic, the most common symptom is the development of an unusually positive view of self, the future, and the world at large. When people are becoming manic, they often find themselves feeling particularly self-confident and experiencing a general sense of well-being. This is often accompanied by beliefs that include overestimating their abilities, an optimism that causes them to view the world as particularly helpful, or an underestimation of the negative consequences of their behaviour. Sometimes this self-confidence can progress to the point where the person's thinking is grandiose and delusional.

When people are becoming manic or hypomanic they may develop a wealth of new ideas and plans that often have the potential to be successful. Unfortunately, when people are manic, it can be difficult to distinguish between good ideas and grandiose delusions. The other thing that often happens during a manic episode is that a person's ability to concentrate and follow through with plans is impaired (particularly when they are overactive or not sleeping much). All of these factors, together with little or no planning, can cause even the best ideas to fail.

During a manic or hypomanic episode, people may forget to consider the potential negative consequences of their behaviour. There is a tendency to rush into an activity or make a decision without carefully considering the pros and cons associated with each choice. They may also be feeling so optimistic that they dismiss the potential negative consequences anyway.

Some people may think they have special powers, particularly in the areas of creativity and personal interactions. These beliefs are maintained when people focus only on events that appear to confirm the belief, while ignoring evidence to the contrary.

Paranoia

Early in the development of mania or hypomania, paranoid thinking is often evident, and may manifest as suspiciousness of others. This suspiciousness is often based on very real events and a history of bad feelings between the person experiencing mania, and the target of his or her paranoia. The paranoia is often maintained by focusing on evidence that confirms the paranoid thoughts. Interactions with the target

person are likely to be tense because of the suspicion involved, and the target person is likely to react in some way to the tension and hostility. This reaction can then be read by the person as confirming their paranoia.

Increased Fluency of Ideas

During mania and hypomania, there is often a flood of new ideas and interests. People with hypomanic symptoms frequently overestimate how much can be accomplished in a given day and underestimate the time it takes to complete tasks. The increase in mental activity that accompanies mania acts to distract people from these tasks they have started doing, making it even more difficult to complete activities.

Thinking Errors in Mania

Social judgement can be impaired in hypomania and is usually quite impaired during manic episodes. People with bipolar disorder often experience a decreased sense of self-awareness during their interactions with others. They may say or do unusual things, or fail to notice the impact that their words or actions have on other people.

On the next page, we will discuss some strategies for preventing a full-blown episode of mania.

The Cognitive Behavioural Approach to Preventing Mania

The cognitive behavioural approach to managing the cognitive and behavioural changes associated with mania or hypomania include:

1. Recognising subtle mood, cognitive, and behavioural changes to allow for early intervention
2. Evaluating and changing distorted or unhelpful thoughts and beliefs, or ideas such as suspiciousness or grandiosity
3. Organising and scheduling of activities, ideas, or plans with the aim of limiting and containing overly-stimulating activities.

In this module, let's look at how you can learn to manage your thoughts to prevent mania. First, you need to learn to recognise what sorts of thoughts you get when you are beginning to get high. When you are in an elevated mood, thoughts that are overly positive might not be helpful to you because they could cause your mood to spiral upwards toward mania. Just like negative thoughts are considered to be unhelpful when you are depressed, thoughts that overly positive are also considered unhelpful when your mood is elevated.

On the next two pages, a worksheet called 'The "Balance" Sheet' is provided, together with an example. This worksheet is for you to record those thoughts that might be overly positive when you are in a mildly or moderately elevated mood. You might not know whether or not they are helpful and that's why it is a good idea to write them down. You might find that some of your thoughts seem a little grandiose and are quite different from how you would normally view a situation, and these would be considered unhelpful to you. The next step is then to develop a more balanced thought in place of the initial unhelpful thought. Here are some questions you can ask yourself to help you develop a balanced thought:

- What evidence do I have that my thoughts are true?
- What facts or details might I have ignored or overlooked?
- What other ways are there of viewing the situation?
- If I were not feeling high, what would I be thinking?
- What is a balanced view of this situation?

It is probably very helpful if you practise this exercise before you start feeling high. Remember that preventing a full-blown episode of mania is all about EARLY recognition of the warning signs and intervening quickly before they escalate. Recognising early warning signs of mania is essential because cognitive behavioural therapy might not be very effective once a person is in a full-blown manic episode.


On the next page is an example of how you can complete the worksheet. Then, turn to page 6 and try out the exercise for yourself!

An Example of a “Balance” Sheet

When I am feeling euphoric, I often think:



An Alternative, Balanced View



Ask yourself:

- What evidence do I have that my thoughts are true?
- What facts or details might I have ignored or overlooked?
- What other explanations could there possibly be?
- Are there other ways of viewing the situation?
- What is a balanced view of this situation?

Wow! I'm feeling really creative and energized right now. Everyone else seems so uninspired!

Hmm... I think my mood might be getting a little elevated. While it feels good to be creative and energized, it might be a good idea to keep a lid on things. Anyway, it's not really fair to say that everyone seems uninspired - they're all working really hard at finishing this project ...

Shopping seems like a really good idea to me! I want a new pair of boots and a new jacket!


Hmm... do I really need those new things? What are my finances like right now? Not all that good, I reckon. Anyway, it'll be spring soon - I might not need a new pair of boots or a new jacket right now...

A "Balance" Sheet

When I am feeling euphoric, I often think:



An Alternative, Balanced View



Ask yourself:

- What evidence do I have that my thoughts are true?
- What facts or details might I have ignored or overlooked?
- What other explanations could there possibly be?
- Are there other ways of viewing the situation?
- What is a balanced view of this situation?

Module Summary

- The symptoms of mania, hypomania, or mixed episodes usually emerge over a period of days and may have a pattern in their development
- Cognitive changes associated with hypomania and mania include increased self-confidence and optimism, and an increase in creative ideas and interests
- Sometimes, paranoid thinking might be present and is demonstrated through suspiciousness of others
- Judgement can sometimes be impaired when mood is elevated
- Just like negative thoughts are considered to be unhelpful when you are depressed, thoughts that overly positive are also considered unhelpful when your mood is elevated.
- It is important to be aware of these thoughts, to write them down and develop a balanced view of the initial thought. The ultimate purpose of this is to prevent a full-blown episode of mania
- Try to practise this exercise before you actually start feeling high
- Preventing a full-blown episode of mania is all about EARLY recognition of the warning signs and intervening quickly before they escalate



Coming up next ...

In the next module, we will discuss behavioural symptoms of hypomania and mania, and identify specific strategies you can use to prevent full-blown episodes.

About The Modules

Contributors

Laura Smith (MPsych¹)
Centre for Clinical Interventions

Dr Louella Lim (DPsych²)
Centre for Clinical Interventions

¹Masters of Psychology (Clinical Psychology)

²Doctor of Psychology (Clinical Psychology)

Background

The concepts and strategies in these modules have been developed from evidence based psychological practice, primarily Cognitive Behaviour Therapy. CBT for bipolar disorder is based on the approach that adjunctive psychological treatment is helpful to improve understanding of the illness, medication adherence, awareness of early warning signs of mood episodes, quality of life and to reduce symptoms.

References

These are some of the professional references that informed the development of modules in this information package.

- Basco, M.R., & Rush, A.J. (2005) Cognitive Behavioral Therapy for Bipolar Disorder (Second Edition). New York: The Guildford Press
- Bauer, M.S. & McBride, L. (2003) Structured Group Psychotherapy for Bipolar Disorder: The Life Goals Program (Second Edition). New York: Springer Publishing Company
- Goodwin, F.K. & Jamison K.R. (1990) Manic Depressive Illness. Oxford; Oxford University Press
- Johnson, S.L. & Leahy, R.L. (2004) Psychological Treatment of Bipolar Disorder. New York: The Guildford Press
- Lam, D.H., Jones, S.H., Hayward, P., & Bright, J.A. (1999) Cognitive Therapy for Bipolar Disorder: A therapist's guide to concepts, methods and practice. Chichester: John Wiley & Sons Ltd
- Miklowitz, D.J. (2001) Bipolar disorder. In David H. Barlow (Ed.) Clinical Handbook of Psychological Disorders – Third Edition (pp 523-561). New York: Guilford Press
- Newman, C.F., Leahy, R.L., Beck, A.T., Reilly-Harrington, N.A., & Gyulai, L. (2002) Bipolar Disorder: A cognitive therapy approach. Washington: American Psychological Association

“KEEPING YOUR BALANCE”

This module forms part of:

Lim, L., & Smith, L. (2003). *Keeping your Balance: Coping with Bipolar Disorder*. Perth, Western Australia: Centre for Clinical Interventions.

ISBN: 9780975198520

Created: March 2003